


## Regular Article

# Associations between cortical thickness and anxious/depressive symptoms differ by the quality of early care

Marta Korom<sup>1</sup> , Nim Tottenham<sup>2</sup>, Emilio A. Valadez<sup>3</sup> and Mary Dozier<sup>1</sup> 

<sup>1</sup>Department of Psychological and Brain Sciences, University of Delaware, Newark, DE, USA; <sup>2</sup>Department of Psychology, Columbia University in the City of New York, New York, NY, USA and <sup>3</sup>Department of Human Development and Quantitative Methodology, University of Maryland, College Park, MD, USA

### Abstract

A variety of childhood experiences can lead to anxious/depressed (A/D) symptoms. The aim of the present study was to explore the brain morphological (cortical thickness and surface area) correlates of A/D symptoms and the extent to which these phenotypes vary depending on the quality of the parenting context in which children develop. Structural magnetic resonance imaging (MRI) scans were acquired on 45 children with Child Protective Services (CPS) involvement due to risk of not receiving adequate care (high-risk group) and 25 children without CPS involvement (low-risk group) (range<sub>age</sub> = 8.08–12.14; *M*<sub>age</sub> = 10.05) to assess cortical thickness (CT) and cortical surface area (SA). A/D symptoms were measured using the Child Behavioral Checklist. The association between A/D symptoms and CT, but not SA, differed by risk status such that high-risk children showed decreasing CT as A/D scores increased, whereas low-risk children showed increasing CT as A/D scores increased. This interaction was specific to CT in prefrontal, frontal, temporal, and parietal cortical regions. The groups had marginally different A/D scores, in the direction of higher risk being associated with lower A/D scores. Results suggest that CT correlates of A/D symptoms are differentially shaped by the quality of early caregiving experiences and should be distinguished between high- and low-risk children.

**Keywords:** anxious/depressive symptoms, caregiving quality, cortical thickness, early adversity, pial surface area

(Received 28 June 2020; accepted 28 June 2021)

Important developmental changes in brain structure occur during middle childhood (Gogtay et al., 2004; Lange, 2012; Marsh, Gerber, & Peterson, 2008; Mills et al., 2016). These changes coincide with the emergence of mental health problems, such as anxiety and depression (Beesdo, Knappe, & Pine, 2009). Structural magnetic resonance imaging (MRI) studies have found cortical thickness (CT) and cortical surface area (SA) alterations that corresponded to clinically significant symptoms (Drevets, Price, & Furey, 2008; Ducharme et al., 2014; Feurer et al., 2020; Newman et al., 2016). However, the direction of change – thinner versus thicker CT or reduced versus increased SA – has been inconsistent across studies, with opposite outcomes across some. A possible confound is that many studies do not assess the nature and quality of parenting experiences and therefore fail to account for the context in which symptoms develop (e.g., see Ducharme et al., 2014; Gold et al., 2017).

The early caregiving environment plays a critical role in children's brain development and susceptibility to mental health problems (Cicchetti, 2016; McLaughlin, Weissman, & Bitrán, 2019). Preliminary evidence suggests that the parenting context

uniquely shapes the neural correlates of anxious symptoms (La Buissonnière-Ariza et al., 2019), despite the fact that distinct childhood experiences can lead to similar presenting symptoms – a phenomenon known as equifinality (Cicchetti & Rogosch, 1996). Of note, La Buissonnière-Ariza et al. (2019) have found that in the context of high versus low harsh parenting, unique functional connectivity (FC) patterns predicted fewer versus higher anxious symptoms. The authors interpreted these findings as adaptations to unique environmental demands that are associated with increased vulnerability to anxiety. Overall, initial steps have been taken to explore the FC patterns associated with anxiety in high- and low-risk parenting context, but little is known about whether the rearing context moderates the morphological correlates of anxious symptoms.

To further our understanding of how the quality of the early parenting context shapes the neurobiological correlates of anxious/depressive (A/D) phenotypes, in this cross-sectional study we explored the CT and SA outcomes of a community sample of children at risk for receiving inadequate care and children without known risk for inadequate care during middle childhood. By directly comparing these groups, this study sought to identify multiple neural pathways leading to A/D symptoms and test the hypothesis that the quality of the early caregiving context, despite social determinants that may differ between groups of children at high versus low risk of experiencing early insensitive care, can distinctly shape the morphological variations seen among children with A/D symptoms.

**Author for Correspondence:** Marta Korom, Wolf Hall 108, 105 The Green, University of Delaware, Newark, DE 19716, USA; E-mail: [mkorom@udel.edu](mailto:mkorom@udel.edu)

**Cite this article:** Korom M, Tottenham N, Valadez EA, Dozier M (2021). Associations between cortical thickness and anxious/depressive symptoms differ by the quality of early care. *Development and Psychopathology* 1–12. <https://doi.org/10.1017/S0954579421000845>

## Early Adversities, Brain Development, and Mental Health Outcomes

The early years of brain development are characterized by profound structural and functional transformations. During the first two years of life the brain doubles in size with regionally heterogeneous developmental patterns (Dobbing & Sands, 1973; Lenroot & Giedd, 2006; Li et al., 2019; Stiles & Jernigan, 2010), and is more sensitive to environmental stimuli than in later years (Tierney & Nelson, 2009). Neural growth occurs in an interpersonal context, where the quality and nature of caregiving practices play a critical role in shaping this change (Cicchetti & Curtis, 2006; Newman, Sivaratnam, & Komiti, 2015). Growing evidence suggests that even normative variability in the quality of caregiving (e.g., parental sensitivity) during infancy is associated with measurable alterations in brain morphology both during infancy (Rifkin-Graboi et al., 2015) and prospectively during middle childhood (Kok et al., 2015). More severe forms of malevolent care, such as abuse or neglect, alter children's structural and functional brain development (Callaghan & Tottenham, 2016b), as indicated by abnormalities in white matter tract integrity (Choi, Jeong, Polcari, Rohan, & Teicher, 2012; Hanson et al., 2013; Huang, Gundapuneedi, & Rao, 2012), altered FC in the fronto-limbic system (Gee et al., 2013; Herringa et al., 2013; Jedd et al., 2015), reduced neuroplasticity (Callaghan & Tottenham, 2016a), increased or atrophied dendritic arborization (Bennett & Diamond, 1996; Bennett, Rosenzweig, Diamond, Morimoto, & Hebert, 1974; Molet et al., 2016), and gray matter reduction (De Brito et al., 2013; Lim, Radua, & Rubia, 2014; Mehta et al., 2009; Tozzi et al., 2020).

Sociodemographic factors, such as low income and low parental educational attainment are also among the risk factors that have been linked to reduced gray matter, estimated total intracranial volume (Brito & Noble, 2014), and altered emotion-regulation neurocircuitry activation (Kim et al., 2013; Liberzon et al., 2015). Although parenting and sociodemographic risk factors have cumulative effects on brain development and associated cognitive outcomes (Chad-Friedman, Botdorf, Riggins, & Dougherty, 2021), randomized clinical trials suggest that the quality of care has unique causal effects on neurodevelopment and associated psychological functioning (Valadez, Tottenham, Tabachnick, & Dozier, 2020). Importantly, the aforementioned morphological alterations, either as a function of insensitive care or sociodemographic determinants, or the cumulative risk of both have been linked to increased vulnerability to psychopathologies, including anxiety and treatment-resistant depression later in development (Busso et al., 2017; Callaghan & Tottenham, 2016a, 2016b; Cicchetti & Toth, 2015; Jensen et al., 2015; Masten & Cicchetti, 2010; Nanni, Uher, & Danese, 2012).

## Cortical Morphology Alterations and Anxious/Depressive Symptoms

There are a few different measures that can assess cortical morphometry. Most studies examine cortical volume, CT, and cortical SA as they relate to psychopathology. Cortical volume is the product of SA and CT, whereas CT and SA are independent of each other both genetically and phenotypically (Grasby et al., 2020; Panizzon et al., 2009; Winkler et al., 2010). In our analyses we will focus on CT and SA.

## Cortical thickness

CT is a measure of cortical gray matter morphology, calculated as the shortest distance between the gray-white matter boundary and the pial surface (Fischl & Dale, 2000). Normative CT maturation is characterized by a linear thinning trajectory across development (LeWinn, Sheridan, Keyes, Hamilton, & McLaughlin, 2017), with increased or reduced thickness associated with anxiety (Gold et al., 2017) and depression (Drevets et al., 2008). Longitudinal studies have also shown that the direction of the association between CT and anxious symptoms may depend on the developmental stage when the MRI assessment occurs (childhood vs. adolescence vs. adulthood) (Ducharme et al., 2014). Such atypical neural signatures indicate that these morphological changes in the geometric organization of the cortex correspond to differences in their functional output (Molet et al., 2016).

Perturbation in CT development is a neurophenotype of non-maltreated, low-risk individuals with A/D symptoms, but reports have been inconsistent in establishing a reliable direction of change. Most studies have found reduced thinning over time (Taylor et al., 2020) or thicker cortices in attention- and control networks (ventromedial prefrontal cortex, orbitofrontal cortex, precentral gyrus, and superior temporal cortices) among a wide age range of low-risk anxious individuals as compared to their nonanxious counterparts (Blackmon et al., 2011; De Bellis et al., 2002; Gold et al., 2017; Schwartz et al., 2012). At the same time, others have identified thinner mean cortices (Newman et al., 2016) or regional specificities in the direction of cortical alterations among adolescents, painting a nuanced picture of the neural correlates of anxious symptoms (Strawn et al., 2013, 2014, 2015). Although these studies are important in examining the effects of anxiety on brain development, they disregard the context in which anxious symptoms develop, which might contribute to the inconsistency of findings across participant pools.

Inadequate care, or in more severe cases, maltreatment, during early childhood is also a predictor of altered CT development and vulnerability to neuropsychiatric disorders later in life (Busso et al., 2017; Kelly et al., 2013). Specifically, both retrospective and prospective reports have identified reductions in medial prefrontal and temporal CT as a predictor of vulnerability to anxious symptoms in children with experiences of caregiving adversities (Busso et al., 2017; Gorka, Hanson, Radtke, & Hariri, 2014).

## Cortical surface area

SA is a measure of both the visible pial surface area and the surface hidden between the sulci (Winkler et al., 2012). SA is closely related to cortical gray matter volume and is a phenotype that has been linked to a variety of psychopathologies (Bois et al., 2015; Prasad et al., 2010; Zhang et al., 2020) and general cognitive ability (Vuoksima et al., 2015). Arguably, SA is a phenotype of great interest in anxiety research, yet existing findings are relatively mixed regarding how SA relates to the quality of early care and severity of anxiety during childhood. For instance, some have reported no main effect of maltreatment on prefrontal cortex SA (Gold et al., 2016), whereas others have shown SA reductions in maltreated versus nonmaltreated children and adolescents in prefrontal (Herzberg et al., 2018; Hodel et al., 2015), temporal, and occipitotemporal cortical regions (Kelly et al., 2013). Reductions in SA among maltreated children are in line with prior work, suggesting that maltreatment is associated with average smaller estimated brain size (Brito & Noble, 2014). The effects

of anxiety on SA are also inconsistent across studies. For instance, in a large group of typically developing participants (age range: 3–20 years), Newman et al. (2016) found a negative association between anxious symptoms and ventromedial prefrontal cortex SA, whereas Bas-Hoogendam et al. (2018) found regional specificities, such that self-reported anxious symptoms negatively correlated with right caudal anterior cingulate cortex and positively with right precuneus at the uncorrected level. Finally, to our knowledge, there is no evidence suggesting that the association between SA and anxiety varies between maltreated and nonmaltreated children.

Other than SA, researchers have studied the effects of anxiety on cortical volume – which is closely related to SA – but most studies focused on young adults and not children. For instance, Rosso et al. (2010) found a positive association between anxiety sensitivity and the volume of the insular cortex among young adults. Qi et al. (2014) found similar results by showing that the positive association between anxious symptoms and cortical volume in the insula was unique to young adults with anxious symptoms as compared to depressed individuals. Overall, the existing literature on cortical SA and volume points toward the prefrontal and insular cortices being potential areas of interest in anxious symptomatology among young adults; however, little is known about these associations among children.

### The Present Study

Taken together, growing evidence suggests that the quality of the early caregiving context can lead to variable morphological outcomes in cortical regions associated with social and emotional processing. These associations are meaningfully related to internalizing symptoms, such as anxiety or depression. In the context of SA, early adversity has been linked to reductions in SA but the findings on how SA relates to anxious symptoms are mixed. With regards to CT, children at risk for maltreatment reliably show thinner cortices in prefrontal, temporal, and parietal regions as their anxiety symptoms increase, whereas nonmaltreated children are more likely to show thicker cortices. However, little is known about whether the caregiving context moderates the association between anxious symptoms and different indices of brain morphology. Preliminary evidence suggests that such moderation effects are present at the functional level (La Buissonnière-Ariza et al., 2019), with high versus low chronic harsh parenting leading to distinct associations between amygdala–rostral anterior cingulate cortex (rACC) FC and anxiety symptoms in adolescents. Overall, initial steps have been taken to explore the FC patterns associated with anxiety in high- and low-risk parenting context, but little is known about whether the rearing context moderates the morphological correlates of anxious symptoms.

The present study addressed several of the limitations of prior research findings contrasting the brain morphological outcomes of a community sample of middle childhood-aged children who received inadequate care (high risk) during infancy versus children without a known history of inadequate care (low risk) with varying severity of A/D symptoms. We hypothesized that children in the high-risk group would show higher anxiety scores and different CT outcomes in frontal and limbic regions than low-risk children with A/D symptoms. We also expected to see smaller SA in regions rendered important in emotion regulation, but we did not have strong predictions about whether SA and anxiety would show unique associations between high- and low-risk caregiving contexts. Characterizing how early experiences shape

morphometric changes during middle childhood could advance our understanding of individual differences and morphological vulnerabilities associated with A/D symptoms and inform research on prevention and psychotherapeutics.

## Method and Materials

### Sample

Participants included 45 high-risk and 25 low-risk children. Families in the high-risk group were referred by Child Protective Services (CPS) from a large mid-Atlantic city as part of a diversion from foster care program for parents deemed at risk for losing their children to foster care. Parents were invited to enroll in a randomized clinical trial testing the efficacy of an early parenting intervention. Families were enrolled in the intervention before the participating children turned 2 years old. The goal of the intervention was to enhance responsive and sensitive caregiving among parents at risk for providing inadequate care to their children. We did not receive access to formal CPS records or information about whether the allegations were substantiated, so we cannot speak to whether children in the high-risk sample had substantiated reports of maltreatment, such as abuse or neglect. However, we know that they were all deemed at risk for not receiving adequate care due to factors such as homelessness, possible maltreatment, parental substance abuse, maltreatment of other children in the family, physical or educational neglect, or domestic violence. Therefore, we characterize these children as high risk.

At the time of the intervention, all but one participating infant were living with their biological mothers. During the middle childhood assessment all but two children were living with their biological parents (the aforementioned child and another high-risk child were placed in the legal custody of their aunts).

At the time of the middle childhood assessment, an age and race-matched group of typically developing 8-year-old children was also recruited in local community centers and schools. Exclusion criteria in this low-risk group included prior history of CPS-involvement, homelessness, or family history of drug abuse at the time of enrollment.

A subset of children from the larger study was invited for an MRI scanning. Inclusion criteria included having an IQ higher than 70 and the absence of neurological disorders. Given that we did not find significant intervention effects on structural brain outcomes, we collapsed the high-risk experimental and control intervention groups into a high-risk group for all analyses.

An initial sample of 80 children underwent MRI scanning. Ten participants' ( $N = 10$ ) data were excluded due to failed data processing ( $N = 5$ ) or poor image quality ( $N = 5$ ; high-risk group excluded data:  $N = 9$ , low-risk group excluded data:  $N = 1$ ). The final sample included 70 children (age range = 8.08–12.14 years,  $M_{\text{age}} = 10.05$  years), of whom 45 were in the high-risk group (25 females) and 25 in the low-risk comparison group (13 females). The high- and low-risk groups did not differ significantly in sex distribution [ $\chi^2(1, N = 70) = .082, p = .78$ ] or age [ $t(68) = -.68, p = .50$ ]. The sample was racially diverse (see more information on racial/ethnic information in Table 1).

### Procedure

Prior to scanning, parents completed the Child Behavioral Checklist (CBCL; Achenbach, Dumenci, & Rescorla, 2003) to

**Table 1.** Sociodemographic and developmental characteristics of high- and low-risk children ( $n = 70$ )

	High-risk group ( $N = 45$ )		Low-risk group ( $N = 25$ )		Group difference	
	<i>M</i>	<i>SE</i>	<i>M</i>	<i>SE</i>	<i>t</i>	<i>p</i> value <sup>a</sup>
Sex, No. (%)	Female, $N = 25$ (62.5%)		Female, $N = 13$ (52%)		$\chi^2 = .082$	.775
	Male, $N = 20$ (37.5%)		Male, $N = 12$ (48%)			
Race/Ethnicity, No. (%)	African American = 31 (68.88%)		African American = 11 (44%)			
	Biracial = 7 (15.55%)		Biracial = 5 (20%)			
	Caucasian = 1 (2.22%)		Caucasian = 7 (28%)			
	Hispanic = 6 (13.33%)		Hispanic = 1 (4%)			
	Did not answer = 0 (0%)		Did not answer = 1 (4%)			
Age (years)						
Age at MRI scan	9.99	.92	10.14	.65	.76	.449
Age at CBCL A/D assessment	9.34	0.57	9.6	0.3	2.118	.038
CBCL – A/D score (sqrt)	0.74	0.75	1.09	0.89	1.770	.081 <sup>b</sup>
Family income (USD)	24,864.71	4,033.56	52,859.2	5,411.59	4.15	.000 <sup>c</sup>
Parental education	2.31	.25	3.72	0.25	5.62	.000 <sup>c</sup>
Brain volume						
Total gray Matter volume (mm <sup>3</sup> )	683,113.54	9,684.29	709,245.47	12,264.07	1.973	.053 <sup>d</sup>
Total intracranial Volume (mm <sup>3</sup> )	1,386,259.05	22,634.35	1,413,119.72	28,663.87	.868	.389 <sup>d</sup>
Cortical gray Matter (mm <sup>3</sup> )	523,004.29	8,319.89	546,988.23	10,536.21	2.108	.039 <sup>d</sup>

A/D = anxious/depressed; CBCL = Child Behavioral Checklist; MRI = magnetic resonance imaging

<sup>a</sup>Significant at the  $p = .05$  level, two-sided test.

<sup>b</sup>Regression model controlled for age and sex.

<sup>c</sup>Educational background was measured on a scale of 6 [1 = did not complete high-school; 2 = GED; 3 = high-school diploma; 4 = some college; 5 = 4-year college degree; 6 = postgraduate degree (MA, MBA, PhD, JD, MD)]

<sup>d</sup>Regression model controlled for age, sex, and the amount of motion in the MRI images.

assess children's anxious/depressive symptomatology and the Brief Symptom Inventory (BSI; Derogatis, 1975) to report on their own mental health status. The average time between the completion of the parent-reported measures and the MRI scanning was 7.29 months ( $SD = 8.5$  months, range 0.7 to 28.67 months). CBCL profiles are relatively stable in pediatric samples, even across a 6- to 8-year period (Biederman et al., 2001; Frizzo, Pedrini, Souza, Bandeira, & Borsa, 2015; Mattison and Spitznagel, 1999; McElroy, Belsky, Carragher, Fearon, & Patalay, 2018; Verhulst and Van der Ende, 1995), which decreases the likelihood that the variability in the time between CBCL completion and the MRI scan in our influenced the results.

Children were accompanied by a female primary caregiver (biological mother for 68, biological aunts for two) to the MRI scan. All children passed the MRI safety screening and had an IQ score of at least 70. Children's comfort in a mock MRI scanner was assessed, followed by the MRI scan. The study protocol was approved by the Institutional Review Board at the University of Delaware.

### Image acquisition and processing

T1-weighted magnetization prepared - rapid gradient echo sequences (MP-RAGE) ( $1 \times 1 \times 1$  mm isometric voxels) were acquired at the Center for Biomedical and Brain Imaging at the University of Delaware using a 3-Tesla Siemens MAGNETOM Prisma<sup>fit</sup> scanner, equipped with a 20-channel head coil for multiband capability. We used standard image processing procedures

in the FreeSurfer image analysis software suite (Version 6, <http://surfer.nmr.mgh.harvard.edu>), including cortical mantle reconstruction and spatial smoothing of 10 mm FWHM. Technical details of the FreeSurfer procedures are described elsewhere (Dale, Fischl, & Sereno, 1999; Fischl et al., 2002, 2004; Fischl & Dale, 2000).

### Quality control of the MRI images

Of the 80 brains, 75 were successfully processed with FreeSurfer, followed by two rounds of quality control. First, a graduate student rated the quality of each image on a scale of 1 through 4 (1 = great segmentation, no motion, include in analyses; 2 = good segmentation, some motion, include in analyses; 3 = satisfactory segmentation, substantial motion, include in analyses; 4 = poor segmentation, extensive motion, exclude from analyses) based on Afacan et al.'s (2016) guidelines. Secondary quality assessment was performed using the Qoala-T supervised-learning tool (Klapwijk, van de Kamp, van der Meulen, Peters, & Wierenga, 2019). Of the 75 processed brains the same five scans were recommended for exclusion by the trained rater and Qoala-T. The low- and high-risk groups did not differ significantly in mean Qoala-T percentile scores ( $t(69) = -.25$ ,  $p = .81$ ) or the rater's scorings [ $t(69) = .48$ ,  $p = .63$ ], and the correlation between the two rating schemes was  $r(68) = -.43$ ,  $p = .00$ . The MRI images were not edited manually to avoid the introduction

190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251

of unsystematic noise. We controlled for image quality, as rated by the trained graduate student, in our analyses.

## Measures

### Demographics

We assessed age, sex, race, family income, caregiver educational background, and caregiver mental health outcomes. Eight families in the high-risk group did not report on their income; therefore, the average indicated income in the high-risk group ( $M = \$34,862.74$ ) represents the income of the 37 families with reported data.

### Anxious/depressive symptoms

#### *Child Behavioral Checklist – anxious/depressed subscale (CBCL A/D)*

The CBCL is a parent-reported standardized questionnaire that reliably assesses symptoms of anxiety and depression between the ages of 6 and 18 (Achenbach & Rescorla, 2001). The internal consistency of the CBCL was excellent in the present sample (overall Cronbach  $\alpha = .946$ ; high-risk group:  $\alpha = .969$ ; low-risk group:  $\alpha = .902$ ). The A/D subscale is a composite of 13 items and has been widely used in a variety of pediatric populations. One item related to suicidal thoughts was removed from the questionnaire, resulting in a 12-item scale in the present study (overall internal consistency in our sample:  $\alpha = .678$ ; high-risk group:  $\alpha = .709$ ; low-risk group:  $\alpha = .628$ ).

#### *Brief Symptom Inventory (BSI)*

The BSI is a 53-item self-report symptom inventory that measures psychiatric symptom patterns on a 5-point Likert scale (0 – *not at all*; 4 – *very much*). It has nine primary symptom dimensions and two global indices of psychological functioning (Derogatis, 1975).

### Statistical analyses

We used linear regression analyses in R (Version 3.5.1; R Core Team, 2019) to examine the main effect of risk status on A/D symptoms. We then examined the association between risk status, A/D symptoms, CT and SA using FreeSurfer's Qdec application ([www.surfer.nmr.mgh.harvard.edu](http://www.surfer.nmr.mgh.harvard.edu)). To test our CT-related hypotheses, we performed vertex-wise general linear model analyses across the entire cortex with an interaction term between A/D symptoms and risk status. The main effect was included in the model. In our SA models, we examined the main effect of risk status and CBCL A/D symptoms, as well as the interaction between the two in separate models. Once again, the main effect was included in the SA model with the interaction term.

We used a Gaussian kernel of 10 mm full width at half maximum (FWHM) to spatially smooth the data. The vertex-wise threshold of significance was  $p < .01$ . We corrected for multiple comparisons using Monte Carlo simulations with a cluster-wide corrected threshold being  $-\log_{10}(p)$  (Hagler, Saygin, & Sereno, 2006). Clusters that survived multiple comparisons corrections were used then as masks to calculate mean CT and SA in that region for each study participant, which then were entered into a regression model and were visualized in R. Only the clusters that survived multiple comparisons correction are reported. All statistical models included sex and age at the time of scanning to follow the example of previous studies on brain morphology (Busso et al., 2017; Gold et al., 2016; McLaughlin, Sheridan, &

Lambert, 2014) and because we hypothesized that anxiety-related alterations in morphology would be unique to the caregiving context and not simply explained by normative, age-related thinning (LeWinn et al., 2017) or age-related changes in anxious symptoms (Beesdo et al., 2009).

In addition, we controlled for image quality (as rated by the trained graduate student). Finally, in separate models, we also controlled for estimated total intracranial volume (ETIV) as a covariate to test if the effects changed with an increase in brain size.

To assess the effect of possible confounding sociodemographic variables on the association between A/D scores and CT, we completed secondary analyses in which we controlled for income, parental education, and race (in addition to sex, age, and image quality). In these models we also controlled for ETIV to make our results comparable to prior studies (e.g., McLaughlin et al., 2014).

## Results

### Demographics

High- and low-risk groups differed in educational background and income, with low-risk parents having significantly higher educational background and income than the high-risk group. In our primary analyses we did not control for parental education and income to avoid biasing the estimate of the risk status, A/D, and interaction effects on CT (Miller & Chapman, 2001). For detailed demographic information, see Table 1.

### Preliminary analyses


Most parents reported subclinical levels of A/D symptoms for their children (range 0–11;  $M = 1.41$ ,  $SD = 1.96$ ), which resulted in positively skewed A/D scores. To adjust for the skew, the square root of the raw A/D scores (range 0–3.32;  $M = .87$ ,  $SD = .81$ ) were used in all analyses. The high-risk and the low-risk groups showed a marginally significant group difference in A/D symptoms [ $b = .36$ ,  $t(66) = 1.77$ ,  $p = .08$ ], in the direction of lower risk being associated with higher anxiety. Age and sex were included in the preliminary analyses as covariates of noninterest.

The parents of high- and low-risk children did not differ in their own report of psychopathology as measured by the nine primary symptom dimensions, the Global Severity Index, and Positive Symptom Total Index of the BSI. Given the null effects on the BSI, we did not control for caregiver mental health outcomes in our models.

### Primary analyses

#### *Risk status and anxious/depressive (A/D) symptom interaction in predicting cortical morphology*

**Cortical thickness.** Our primary hypotheses aimed to test interaction effects between risk status and A/D symptoms in predicting CT. See Figure 1 for visual representation of our significant findings. In the right hemisphere, three clusters (clusters A, B, and C) survived Monte Carlo multiple comparison corrections. These clusters peaks were in the (A) precentral, (B) superior parietal, and (C) pericalcarine cortices. In the left hemisphere, six clusters emerged (clusters D through I) with their peaks in the (D) precentral-, (E and I) superior frontal-, (F) pericalcarine-, (G) superior parietal-, and (H) transverse temporal cortices. For



**Figure 1.** Significant risk status by Child Behavioral Checklist (CBCL) anxious/depressed (A/D) symptoms interactions on cortical thickness (CT) on the group average brain, following Monte Carlo multiple comparison corrections in both hemispheres. Greater A/D symptoms were associated with decreasing CT among the high-risk group and increasing CT in the low-risk group. The model controlled for age, sex, and image quality. The scatter plots are prototypical associations that were similar across all the clusters. RH – right hemisphere; LH – left hemisphere; A – RH precentral Cortex (including areas of the superior frontal-, paracentral-, caudal middle frontal-, rostral middle frontal-, pars triangularis-, rostral middle frontal-, pars opercularis-, and postcentral cortex); B – RH superior parietal cortex (including areas of the precuneus, lateral occipital-, and inferior parietal cortex); C – RH pericalcarine Cortex (including areas of the lateral occipital-, lingual and fusiform cortex); D – LH precentral Cortex (including areas of the caudal middle frontal-, and rostral middle frontal cortex); E – LH superior frontal Cortex (including areas of rostral middle frontal, medial orbitofrontal, rostral anterior cingulate, and caudal anterior cingulate cortex); F – LH pericalcarine Cortex (including areas lingual and lateral occipital cortex); G – LH superior parietal Cortex (including areas of the postcentral-, inferior parietal-, and lateral occipital cortex); H – LH transverse temporal Cortex (including areas of the insula, superior temporal and transverse temporal cortex); I – LH superior frontal Cortex (including areas of the paracentral-, precuneus, and posterior cingulate cortex).

**Table 2.** Left and right hemisphere cluster characteristics for the clusters that survived multiple comparisons correction

	Cluster size (mm <sup>2</sup> )	Peak within cluster <i>t</i>	Approximate coordinates in MNI space ( <i>x</i> , <i>y</i> , <i>z</i> )		
			<i>x</i>	<i>y</i>	<i>z</i>
Left Hemisphere/cortical thickness/interaction effects					
Superior parietal cortex	2,823.44	-3.366	-22.2	-53.5	40.8
Pericalcarine cortex	2,631.15	-3.986	-7.8	-84.1	4.2
Precentral cortex	2,484.08	-3.907	-34.4	-6.8	44.3
Superior frontal cortex A	1,481.51	-3.365	-10.1	-6.7	45
Superior frontal cortex B	1,333.03	-3.959	-10.9	42	6
Transverse temporal cortex	1,237.46	-2.897	-43.9	-20.5	5.5
Right hemisphere/cortical thickness/interaction effects					
Precentral cortex	6,693.88	-3.944	26.5	-11.2	54.3
Superior parietal cortex	2,774.85	-2.633	24.6	-68.8	37.5
Pericalcarine cortex	1,165.26	-2.209	15.4	-90.3	3.7
Right hemisphere/pial surface area/main effect of risk status					
Caudal middle frontal	1,208.48	4.139	34.1	12.1	28.0
Right hemisphere/pial surface area/main effect of CBCL A/D score					
Insula	1,592.08	3.828	36.1	-10.8	-2.8

A/D = anxious/depressed; CBCL = Child Behavioral Checklist; anxious/depressed; MNI = Montreal Neurological Institute. All analyses controlled for age, sex, and amount of motion in individual MRI images.

information on cluster characteristics (size, Montreal Neurological Institute [MNI] coordinates and peak), see Table 2. In all the clusters, greater A/D symptoms were associated with decreased CT

among the high-risk children and increased CT in low-risk children (Figures 1). As can be seen in the right panel of Figure 1, as the number of A/D symptoms increase, CT decreases in the high-

**Table 3.** Simple slopes results for the interactions between anxious/depressive outcomes and cortical thickness in the high- and the low-risk group

	High-risk group (N = 45)				Low-risk group (N = 25)				Risk * CBCL A/D score interaction effect			
	Intercept	b	SE	t	Intercept	b	SE	t	b	SE	t	
Left hemisphere												
Superior parietal cortex	2.490	-0.09	.020	-4.474	2.366	.065	.023	2.829	.156	.031	5.085	
Pericalcarine cortex	2.362	-.082	.022	-3.754	2.201	.080	.025	3.207	.163	.033	4.893	
Precentral cortex	3.227	-.101	.024	-4.202	3.079	.064	.027	2.340	.166	.037	4.538	
Superior frontal cortex A	2.912	-.079	.025	-3.108	2.739	.063	.028	2.204	.143	.039	3.646	
Superior frontal cortex B	3.568	-.135	.035	-3.807	3.390	.083	.040	2.068	.218	.054	4.001	
Transverse temporal cortex	3.519	-.116	.033	-3.500	3.359	.060	.038	1.589	.177	.050	3.509	
Right hemisphere												
Precentral cortex	3.122	-.097	.026	-3.676	2.924	.090	.030	3.019	.187	.040	4.700	
Pericalcarine cortex	2.215	-.071	.023	-3.143	2.068	.048	.026	1.860	.119	.034	3.477	
Superior parietal cortex	2.85	-.097	.024	-3.961	2.748	.047	.028	1.688	.144	.037	3.888	

Note. The estimates reflect unstandardized regression coefficients. A/D = anxious/depressed; CBCL = Child Behavioral Checklist

risk group. In contrast, in the right panel, as the number of A/D symptoms increase, CT increases in the low-risk group. For simple slope effects, see Table 3.

**Cortical surface area.** We hypothesized that children in the high-risk group would show smaller cortical areas in regions rendered key for healthy social and emotional functioning. We also tested for significant associations between SA and CBCL A/D scores across the pial surface. Finally, we examined the effect of the interaction term between A/D symptoms and risk status on SA. See Figure 2 for our results. Children in the high-risk group showed significantly smaller SA than low-risk children in the prefrontal cortex, with the cluster peak being in the rostral middle frontal cortex in the right hemisphere (see Figure 2A and Table 4). SA also significantly correlated with CBCL A/D scores in the right hemisphere insular cortex, such that as anxious symptoms increased, SA also increased (see Figure 2B). The effect remained significant even after we controlled for risk status. Finally, risk status did not interact with anxious symptoms in predicting SA.

### Secondary analyses


Our secondary analyses examined if sociodemographic variables, including income, parental education, and race would change the results of our primary analyses. The findings related to CT and SA remained significant. Finally, we tested the additional effect of ETIV on our results. The CT findings and the main effect of anxiety on insular cortex SA remained significant with the inclusion of ETIV in the model; however, the main effect of risk status on SA became nonsignificant.

### Discussion

The present study investigated the association between two indices of cortical morphology, namely CT and CSA, and associated A/D symptoms in high- versus low-risk caregiving environments during middle childhood. We found that in the high-risk group, as the severity of A/D symptoms increased, CT decreased,

whereas in the low-risk group, as A/D symptoms increased, CT increased. We also found a significant main effect of risk status on SA, with children in the high-risk group showing smaller SA in a subregion of the prefrontal cortex than low-risk children. In addition, SA was positively associated with CBCL A/D symptoms in the insular cortex. These associations were observed in cortical regions responsible for higher-level cognitive, and attentional processes and regulation of emotional stimuli (e.g., frontal, medial prefrontal, temporal, and parietal areas). These preliminary findings highlight the importance of assessing the quality of children's caregiving environment when examining the CT correlates of A/D symptoms; however, the results should be viewed in light of the small sample size and the exploratory nature of our results that need future replication.

This exploratory study encourages future studies to consider early caregiving as a moderator of brain-behavior associations, but we cannot tell why we are seeing distinct associations between the two. There can be many explanations as to why distinct morphometric phenotypes were observed in high- and low-risk contexts, all of which are speculative given the present data. For instance, considering that the cortex shows a normative linear thinning trajectory across development, thinner cortices as a function of higher anxious symptoms might reflect accelerated thinning in high-risk contexts. Perhaps chronic experiences of threat and insensitive care in the high-risk group lead to premature maturation of the optimally slow-paced development of the prefrontal- and orbitofrontal cortices in the service of improved self-regulation early in life (Callaghan & Tottenham, 2016b), which might become a vulnerability factor during middle childhood when internalizing symptoms commonly arise. Exposure to challenges can also alter stress hormone production that has the potential to shape brain morphology (Kim & Yoon, 1998; Lupien et al., 1998; McEwen, Nasca, & Gray, 2016; Sapolsky, 1996) by reducing dendritic arborization and soma size, hence leading to reductions in gray matter volume and thickness. In sum, although thinner cortices may be adaptive and may aid high-risk children's emotion regulation in coping with challenges, they likely become a vulnerability factor at an older age in the context of chronic stressors.



**Figure 2.** Significant pial surface area outcomes. 2A – significant main effect of risk status on right hemisphere caudal middle frontal region surface area on the inflated brain surface and the distribution plot of individual average surface area outcomes (the error bars represent  $\pm 3$  standard deviations). 2B – significant main effect of Child Behavioral Checklist (CBCL) anxious/depressed (A/D) scores on insular cortex SA on the inflated brain surface. The scatter plot depicts the positive association between the CBCL A/D scores and SA. Both models controlled for age, sex, and image quality.

**Table 4.** Significant main effects of Child Behavioral Checklist (CBCL) anxious/depressed (A/D) symptoms and risk status on pial surface area

	Intercept	<i>b</i>	<i>SE</i>	<i>t</i>
Main effect of CBCL A/D on RH SA	.379	.029	.006	4.377
Main effect of risk status on RH SA (high risk < low risk)	1.014	.099	.047	2.114

Note. The estimates reflect unstandardized regression coefficients. RH = right hemisphere; SA = pial surface area.

In the case of children from low-risk backgrounds with A/D symptoms, the significance of altered cortical development, as manifested by thicker cortices, is still debated. Findings have suggested that there is a delay in structural maturation of the medial prefrontal cortex (mPFC) (Ducharme et al., 2014) as well as the white matter tracts between the PFC and limbic structures among anxious adolescents (Miller et al., 2012; Paus, Keshavan, & Giedd, 2008). Thicker cortices might also indicate a compensatory mechanism, where the PFC becomes more active as it tries to control the overly active amygdala (Herringa et al., 2016; McLaughlin, Peverill, Gold, Alves, & Sheridan, 2015). Increased PFC activation can lead to increased cell numbers and myelination, which in turn can result in thicker cortices (Hegarty et al., 2012; Vidal-Pineiro et al., 2020). Finally, dispositional factors, such as genetic vulnerability or temperament, might also shape neural correlates of A/D symptomatology.

Although it is speculative, the observed cortical phenotype among children from low-risk environments are perhaps also related to characteristics of their caregiving environments. Parents of anxious children tend to model anxious behaviors, encourage maladaptive

coping, and engage in overprotective and overcontrolling behaviors (Kendall, 2012). Under such circumstances, the normative maturation of the mPFC may be delayed – as seen in thicker cortices with the increase of anxious symptoms – leading to prolonged dependence on the caregiver for co-regulation, later onset of normative cortical thinning, and poor self-regulation during middle childhood.

Our findings might also be in line with La Buissonnière-Ariza and colleagues' work (2019) where the quality of parenting (high vs. low harsh parenting) predicted unique associations between anxious symptoms and FC patterns between the rACC and the amygdala. Future studies are necessary to confirm this, but the unique morphological profiles observed in our study might be related to distinct FC patterns between the amygdala and the rACC.

We note that children from low-risk and high-risk groups showed a marginally significant difference in their parent-reported anxious symptom severity. More specifically, caregivers of children in the high-risk group were marginally more likely to report fewer anxious symptoms than children in the low-risk group. It is also important to highlight that although early caregiving adversities in general increase children's vulnerability to internalizing psychopathologies, null findings between high- and low-risk children have emerged in prior studies as well (Cicchetti, Rogosch, Gunnar, & Toth, 2010; Dubois-Comtois, Moss, Cyr, & Pascuzzo, 2013; Keiley, Howe, Dodge, Bates, & Petti, 2001; Wiik et al., 2011).

Unlike CT, risk status did not predict unique associations between anxious symptoms and SA. Instead, we found group differences in average SA in the prefrontal cortex, such that children at risk for inadequate care showed smaller SA than low-risk children. These effects are in line with our findings related to total gray matter volume where children in the high-risk group showed reductions compared to the low-risk children (Brito & Noble,

438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499



2014). Given that gray matter volume is closely related to SA (more so than CT) (Winkler et al., 2010), such a group difference was expected.

We also found a significant positive association between SA and CBCL A/D scores in the insular cortex that was unrelated to the quality of early care. Considering that the insular cortex is a central node of the fear circuitry and is a cross-modal integration hub where interoceptive, emotional, sensory, cognitive, and motivational input converge to shape behavior (Gogolla, 2017; Kurth, Zilles, Fox, Laird, & Eickhoff, 2010), our results might indicate increased engagement of the insular cortex in the service of modulation of subclinical anxious symptoms during middle childhood. To our knowledge, insular SA has not been directly linked to anxious symptoms in children, but some studies have found similar associations between insular cortex volume and anxious symptom severity (Rosso et al., 2010). Our results are also in line with a vast literature on the insular cortex showing increased activation among anxious individuals (Kurth et al., 2010), which has the potential to increase insular volume and perhaps SA. Given that our sample showed mainly subclinical levels of anxiety, future studies are encouraged to examine these associations in clinically anxious pediatric populations.

The present study has possible preventive and psychotherapeutic implications, especially in the context of CT. Given that distinct neurophenotypes were associated with anxious symptoms, children from high- and low-caregiving contexts might benefit from different forms of interventions. At the subclinical symptom level, children at risk for developing anxiety may benefit from school-based prevention and early intervention programs. Most of these programs use techniques from cognitive behavioral therapy (CBT), mindfulness-based activities, and psychoeducation and have shown to reduce short- and longer-term anxiety (Neil & Christensen, 2009). Children with clinically significant symptoms may benefit from targeted mindfulness-based interventions (Hofmann, Sawyer, Witt, & Oh, 2010), CBT (Seligman & Ollendick, 2011), and the combination of the two (mindfulness-based CBT; Semple & Lee, 2007), all of which have been identified as powerful treatments for anxiety. To our knowledge, there is no evidence from randomized clinical trials deeming one approach superior to the other for children from high-versus low-risk caregiving contexts. The causal effects of evidence-based interventions on changes in CT are also unknown. A recent study has found that thinner cortices in parietal and occipital regions, including the lingual gyrus, were predictive of worse treatment outcomes among anxious youth (Gold et al., 2017); this study, however, did not assess the quality of the early caregiving context and was not a randomized clinical trial. Nevertheless, if thinner cortices are indicative of worse treatment outcomes in general, thicker cortices among low-risk anxious children might suggest better odds for symptom reduction following therapy than high-risk anxious children with thinner cortices. Although the biological processes underlying these hypothesized mechanisms are speculative, perhaps thicker cortices are indicative of enhanced neuroplasticity. Future studies are needed to address this gap in the literature and systematically investigate the causal effects of therapy on CT development in high- and low-risk contexts to maximize children's therapeutic gains.

### Strengths and Limitations

There are several limitations that need to be addressed. First, the high- and low-risk groups are unbalanced on several sociodemographic factors (e.g., income, parental education, distribution of race). High sociodemographic risk factors tend to coincide

(Smith et al., 2015) and are associated with gray matter volume reductions in high-risk populations (Brito & Noble, 2014; De Bellis et al., 2002; Lim et al., 2018; VanTieghem et al., 2021). Although similar sample characteristics are common in other studies comparing high- versus low-risk groups of children (e.g., see Busso et al., 2017), such group differences might bias or even introduce alternative interpretations of the results. To address this, we statistically accounted for this unbalance in our data and our results did not change, suggesting that risk status continued to predict unique morphological variations over and above socio-demographic determinants. Second, we have limited information about the allegations and the associated parental care that resulted in the families' CPS involvement. Some of the allegations were likely substantiated and were associated with insensitive care, however, they were not severe enough to result in the child's removal from the family. Despite evidence suggesting that both substantiated and unsubstantiated allegations of maltreatment have a similar negative effect on developmental outcomes (Hussey et al., 2005), further research needs to consider and characterize how the type, severity, and frequency of maltreatment relate to mental health outcomes and alterations in brain morphology. Third, the sample size of the low-risk group is relatively small. Underpowered studies are vulnerable to Type 1 errors, therefore studies with a larger sample size are encouraged to replicate our preliminary findings. The present study was exploratory, nevertheless, relative to other studies with a much wider age range, the present study represents a valuable initial step toward characterizing structural changes as a function of A/D symptoms and risk status during a narrow but critical developmental period when mood disorders commonly arise. For more information on age-related changes in CT and anxiety, see Ducharme et al. (2014). In addition, we used a parent-report measure to assess A/D symptoms, which could be influenced by parental biases in the perception and reporting of behavior problems. Despite this limitation, the CBCL is a widely used measure and gives an important assessment of children's mental health problems from the parents' perspective. Lastly, although studying how subclinical levels of A/D symptoms shape the neural phenotypes of children is critical, extrapolations to potential neural changes in pediatric populations with more severe psychopathology remain speculative.

The current study presents preliminary evidence for the effect of the early caregiving environment on the morphological associations of subclinical anxiety symptoms during middle childhood. Future studies are needed to continue to examine the factors influencing the direction of CT differences, as well as explore prospective associations between early-life caregiving experiences and patterns of structural brain development, with the goal to explain individual variability in negative mental health outcomes and the biological characteristics influencing children's therapeutic gains.

**Acknowledgments.** Marta Korom completed data processing, quality control of all images, statistical analyses and wrote the manuscript. As a co-Investigator, Dr. Tottenham helped develop the research design and assisted in analyses and manuscript development. Emilio A. Valadez was responsible for data collection, helped develop the research design, and assisted in manuscript editing. As the primary investigator, Dr. Dozier wrote the grant that funded the study, developed the research design, and assisted in analyses and manuscript development.

**Funding Statement.** This research was supported by the National Institute of Mental Health R01 R01MH074374-07S1 grant.

**Conflicts of Interest.** None.

## References

- Achenbach, T. M., Dumenci, L., & Rescorla, L. A. (2003). DSM-oriented and empirically based approaches to constructing scales from the same item pools. *Journal of Clinical Child & Adolescent Psychology, 32*, 328–340.
- Achenbach, T. M., & Rescorla, L. A. (2001). *Manual for the ASEBA school-Age forms & profiles*. Burlington, VT: University of Vermont, Research Center for Children, Youth, & Families.
- Afacan, O., Erem, B., Roby, D. P., Roth, N., Roth, A., Prabhu, S. P., & Warfield, S. K. (2016). Evaluation of motion and its effect on brain magnetic resonance image quality in children. *Pediatric Radiology, 46*, 1728–1735.
- Bas-Hoogendam, J. M., van Steenbergen, H., Tisseries, R., Houwing-Duistermaat, J. J., Westenberg, P. M., & van der Wee, N. (2018). Subcortical brain volumes, cortical thickness and cortical surface area in families genetically enriched for social anxiety disorder - A multiplex multigenerational neuroimaging study. *EBioMedicine, 36*, 410–428. doi:10.1016/j.ebiom.2018.08.048
- Beesdo, K., Knappe, S., & Pine, D. S. (2009). Anxiety and anxiety disorders in children and adolescents: Developmental issues and implications for DSM-V. *Psychiatric Clinics of North America, 32*, 483–524.
- Bennett, L., & Diamond, C. (1996). Chemical and anatomical plasticity of brain. 1964 [classical article]. *The Journal of Neuropsychiatry and Clinical Neurosciences, 8*, 459–470.
- Bennett, E. L., Rosenzweig, M. R., Diamond, M. C., Morimoto, H., & Hebert, M. (1974). Effects of successive environments on brain measures. *Physiology and Behavior, 12*, 621–631.
- Biederman, J., Monuteaux, M. C., Greene, R. W., Braaten, E., Doyle, A. E., & Faraone, S. V. (2001). Long-term stability of the child behavior checklist in a clinical sample of youth with attention deficit hyperactivity disorder. *Journal of Clinical Child Psychology, 30*, 492–502.
- Blackmon, K., Barr, W. B., Carlson, C., Devinsky, O., DuBois, J., Pogash, D., ... Thesen, T. (2011). Structural evidence for involvement of a left amygdala-orbitofrontal network in subclinical anxiety. *Psychiatry Research: Neuroimaging, 194*, 296–303.
- Bois, C., Ronan, L., Levita, L., Whalley, H. C., Giles, S., McIntosh, A. M., ... Lawrie, S. M. (2015). Cortical surface area differentiates familial high risk individuals who go on to develop schizophrenia. *Biological Psychiatry, 78*, 413–420. doi:10.1016/j.biopsych.2014.12.030
- Brito, N. H., & Noble, K. G. (2014). Socioeconomic status and structural brain development. *Frontiers in Neuroscience, 8*, 276. doi:10.3389/fnins.2014.00276
- Busso, D. S., McLaughlin, K. A., Brueck, S., Peverill, M., Gold, A. L., & Sheridan, M. A. (2017). Child abuse, neural structure, and adolescent psychopathology: A longitudinal study. *Journal of the American Academy of Child and Adolescent Psychiatry, 56*(4), 321–328.e1.
- Callaghan, B. L., & Tottenham, N. (2016a). The neuro-environmental loop of plasticity: A cross-species analysis of parental effects on emotion circuitry development following typical and adverse caregiving. *Neuropsychopharmacology, 41*, 163–176.
- Callaghan, B. L., & Tottenham, N. (2016b). The stress acceleration hypothesis: Effects of early-life adversity on emotion circuits and behavior. *Current Opinion in Behavioral Sciences, 7*, 76–81.
- Chad-Friedman, E., Botdorf, M., Riggins, T., & Dougherty, L. R. (2021). Early childhood cumulative risk is associated with decreased global brain measures, cortical thickness, and cognitive functioning in school-age children. *Developmental Psychobiology, 63*, 192–205. doi:10.1002/dev.21956
- Choi, J., Jeong, B., Polcari, A., Rohan, M. L., & Teicher, M. H. (2012). Reduced fractional anisotropy in the visual limbic pathway of young adults witnessing domestic violence in childhood. *NeuroImage, 59*, 1071–1079.
- Cicchetti, D. (2016). Socioemotional, personality, and biological development: Illustrations from a multilevel developmental psychopathology perspective on child maltreatment. *Annual Review of Psychology, 67*, 187–211. doi: 10.1146/annurev-psych-122414-033259
- Cicchetti, D., & Curtis, W. J. (2006). The developing brain and neural plasticity: Implications for normality, psychopathology, and resilience: Developmental neurosciences. In D. Cicchetti & D. J. Cohen (Eds.), *Developmental psychopathology: Developmental neuroscience* (2nd ed., Vol. 2, pp. 1–64). Hoboken, NJ: John Wiley & Sons, Inc.
- Cicchetti, D., & Rogosch, F. A. (1996). Equifinality and multifinality in developmental psychopathology. *Development and Psychopathology, 8*, 597–600.
- Cicchetti, D., Rogosch, F. A., Gunnar, M. R., & Toth, S. L. (2010). The differential impacts of early physical and sexual abuse and internalizing problems on daytime cortisol rhythm in school-aged children. *Child Development, 81*, 252–269.
- Cicchetti, D., & Toth, S. L. (2015). Child maltreatment. In R. M. Lerner & M. E. Lamb (Eds.), *Handbook of child psychology and developmental science* (Vol. 3, pp. 513–563). Hoboken, NJ: John Wiley & Sons, Inc.
- Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. *NeuroImage, 9*, 179–194.
- De Bellis, M. D., Keshavan, M. S., Shifflett, H., Iyengar, S., Dahl, R. E., Axelson, D. A., ... Ryan, N. D. (2002). Superior temporal gyrus volumes in pediatric generalized anxiety disorder. *Biological Psychiatry, 51*, 553–562.
- De Brito, S. A., Viding, E., Sebastian, C. L., Kelly, P. A., Mechelli, A., Maris, H., & McCrory, E. J. (2013). Reduced orbitofrontal and temporal grey matter in a community sample of maltreated children. *Journal of Child Psychology and Psychiatry and Allied Disciplines, 54*, 105–112.
- Derogatis, L. R. (1975). *Brief symptom inventory*. Baltimore: Clinical Psychometric Research.
- Dobbing, J., & Sands, J. (1973). Quantitative growth and development of human brain. *Archives of Disease in Childhood, 48*, 757–767.
- Drevets, W. C., Price, J. L., & Furey, M. L. (2008). Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Structure and Function, 213*, 93–118.
- Dubois-Comtois, K., Moss, E., Cyr, C., & Pascuzzo, K. (2013). Behavior problems in middle childhood: The predictive role of maternal distress, child attachment, and mother-child interactions. *Journal of Abnormal Child Psychology, 41*, 1311–1324.
- Ducharme, S., Albaugh, M. D., Hudziak, J. J., Botteron, K. N., Nguyen, T. V., Truong, C., ... Karama, S. (2014). Anxious/depressed symptoms are linked to right ventromedial prefrontal cortical thickness maturation in healthy children and young adults. *Cerebral Cortex, 24*, 2941–2950.
- Feurer, C., Suor, J. H., Jimmy, J., Klumpp, H., Monk, C. S., Phan, K. L., & Burkhouse, K. L. (2020). Differences in cortical thinning across development among individuals with and without anxiety disorders. *Depression and Anxiety, 38*, 372–381. doi:10.1002/da.23096
- Fischl, B., & Dale, A. M. (2000). Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of the National Academy of Sciences of the United States of America, 97*, 11050–11055.
- Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., ... Dale, A. M. (2002). Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron, 33*, 341–355.
- Fischl, B., Van Der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D. H., ... Dale, A. M. (2004). Automatically parcellating the human cerebral cortex. *Cerebral Cortex, 14*, 11–22.
- Frizzo, G. B., Pedrini, J. R., Souza, D. S., Bandeira, D. R., & Borsa, J. C. (2015). Reliability of child behavior checklist and teacher's report form in a sample of Brazilian children. *Universitas Psychologica, 14*, 149–156. doi:10.11144/Javeriana.upsy14-1.rcbc
- Gee, D. G., Humphreys, K. L., Flannery, J., Goff, B., Telzer, E. H., Shapiro, M., ... Tottenham, N. (2013). A developmental shift from positive to negative connectivity in human amygdala-prefrontal circuitry. *Journal of Neuroscience, 33*, 4584–4593.
- Gogolla, N. (2017). The insular cortex. *Current Biology, 27*, R580–R586. doi:10.1016/j.cub.2017.05.010
- Gogtay, N., Giedd, J. N., Lusk, L., Hayashi, K. M., Greenstein, D., Vaituzis, A. C., ... Thompson, P. M. (2004). Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences, 101*, 8174–8179.
- Gold, A. L., Sheridan, M. A., Peverill, M., Busso, D. S., Lambert, H. K., Alves, S., ... McLaughlin, K. A. (2016). Childhood abuse and reduced cortical thickness in brain regions involved in emotional processing. *Journal of Child Psychology and Psychiatry, 57*, 1154–1164.
- Gold, A. L., Steuber, E. R., White, L. K., Pacheco, J., Sachs, J. F., Pagliaccio, D., ... Pine, D. S. (2017). Cortical thickness and subcortical gray matter volume in pediatric anxiety disorders. *Neuropsychopharmacology, 42*, 2423–2433.

- Gorka, A. X., Hanson, J. L., Radtke, S. R., & Hariri, A. R. (2014). Reduced hippocampal and medial prefrontal gray matter mediate the association between reported childhood maltreatment and trait anxiety in adulthood and predict sensitivity to future life stress. *Biology of Mood and Anxiety Disorders*, 4, 1–10.
- Grasby, K. L., Jahanshad, N., Painter, J. N., Colodro-Conde, L., Bralten, J., Hibar, D. P., ... Amlien, I. K. ... Enhancing NeuroImaging Genetics through Meta-Analysis Consortium (ENIGMA) – Genetics working group (2020). The genetic architecture of the human cerebral cortex. *Science*, 367, eaay6690. doi:10.1126/science.aay6690
- Hagler, D. J., Saygin, A. P., & Sereno, M. I. (2006). Smoothing and cluster thresholding for cortical surface-based group analysis of fMRI data. *NeuroImage*, 33, 1093–1103.
- Hanson, J. L., Adluru, N., Chung, M. K., Alexander, A. L., Davidson, R. J., & Pollak, S. D. (2013). Early neglect is associated with alterations in white matter integrity and cognitive functioning. *Child Development*, 84, 1566–1578.
- Hegarty, C. E., Foland-Ross, L. C., Narr, K. L., Townsend, J. D., Bookheimer, S. Y., Thompson, P. M., & Altschuler, L. L. (2012). Anterior cingulate activation relates to local cortical thickness. *NeuroReport*, 23, 420–424.
- Herrington, R. J., Birn, R. M., Ruttle, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., & Essex, M. J. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 19119–19124.
- Herrington, R. J., Burghy, C. A., Stodola, D. E., Fox, M. E., Davidson, R. J., & Essex, M. J. (2016). Enhanced prefrontal-amygdala connectivity following childhood adversity as a protective mechanism against internalizing in adolescence. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, 1, 326–334. doi:10.1016/j.bpsc.2016.03.003
- Herzberg, M. P., Hodel, A. S., Cowell, R. A., Hunt, R. H., Gunnar, M. R., & Thomas, K. M. (2018). Risk taking, decision-making, and brain volume in youth adopted internationally from institutional care. *Neuropsychologia*, 119, 262–270. doi:10.1016/j.neuropsychologia.2018.08.022
- Hodel, A. S., Hunt, R. H., Cowell, R. A., Van Den Heuvel, S. E., Gunnar, M. R., & Thomas, K. M. (2015). Duration of early adversity and structural brain development in post-institutionalized adolescents. *NeuroImage*, 105, 112–119. doi:10.1016/j.neuroimage.2014.10.020
- Hofmann, S. G., Sawyer, A. T., Witt, A. A., & Oh, D. (2010). The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of Consulting and Clinical Psychology*, 78, 169–183.
- Huang, H., Gundapuneedi, T., & Rao, U. (2012). White matter disruptions in adolescents exposed to childhood maltreatment and vulnerability to psychopathology. *Neuropsychopharmacology*, 37, 2693–2701.
- Hussey, J. M., Marshall, J. M., English, D. J., Knight, E. D., Lau, A. S., Dubowitz, H., & Kotch, J. B. (2005). Defining maltreatment according to substantiation: Distinction without a difference? *Child Abuse & Neglect*, 29, 479–492. doi:10.1016/j.chiabu.2003.12.005
- Jedd, K., Hunt, R. H., Cicchetti, D., Hunt, E., Cowell, R. A., Rogosch, F. A., ... Thomas, K. M. (2015). Long-term consequences of childhood maltreatment: Altered amygdala functional connectivity. *Development and Psychopathology*, 27, 1577–1589.
- Jensen, S. K., Dickie, E. W., Schwartz, D. H., Evans, C. J., Dumontheil, I., Paus, T., & Barker, E. D. (2015). Effect of early adversity and childhood internalizing symptoms on brain structure in young men. *JAMA Pediatrics*, 169, 938–946. doi:10.1001/jamapediatrics.2015.1486
- Keiley, M. K., Howe, T. R., Dodge, K. A., Bates, J. E., & Pettit, G. S. (2001). The timing of child physical maltreatment: A cross-domain growth analysis of impact on adolescent externalizing and internalizing problems. *Development and Psychopathology*, 13, 891–912.
- Kelly, P. A., Viding, E., Wallace, G. L., Schaer, M., De Brito, S. A., Robustelli, B., & McCrory, E. J. (2013). Cortical thickness, surface area, and gyrification abnormalities in children exposed to maltreatment: Neural markers of vulnerability? *Biological Psychiatry*, 74, 845–852.
- Kendall, P. C. (2012). Anxiety disorders in youth. In P. C. Kendall (Ed.), *Child and adolescent therapy: Cognitive-behavioral procedures* (pp. 143–189). New York, NY, US: Guilford Press.
- Kim, P., Evans, G. W., Angstadt, M., Ho, S. S., Sripada, C. S., Swain, J. E., ... Phan, K. L. (2013). Effects of childhood poverty and chronic stress on emotion regulatory brain function in adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 18442–18447. doi:10.1073/pnas.1308240110
- Kim, J. J., & Yoon, K. S. (1998). Stress: Metaplastic effects in the hippocampus. *Trends in Neuroscience*, 21, 505–509.
- Klapwijk, E. T., van de Kamp, F., van der Meulen, M., Peters, S., & Wierenga, L. M. (2019). Qoala-T: A supervised-learning tool for quality control of FreeSurfer segmented MRI data. *NeuroImage*, 189, 116–129.
- Kok, R., Thijssen, S., Bakermans-Kranenburg, M. J., Jaddoe, V. W., Verhulst, F. C., White, T., ... Tiemeier, H. (2015). Normal variation in early parental sensitivity predicts child structural brain development. *Journal of the American Academy of Child and Adolescent Psychiatry*, 54, 824–831.e1. doi:10.1016/j.jaac.2015.07.009
- Kurth, F., Zilles, K., Fox, P. T., Laird, A. R., & Eickhoff, S. B. (2010). A link between the systems: Functional differentiation and integration within the human insula revealed by meta-analysis. *Brain Structure and Function*, 214, 519–534. doi: 10.1007/s00429-010-0255-z
- La Buissonnière-Ariza, V., Séguin, J. R., Nassim, M., Boivin, M., Pine, D. S., Lepore, F., ... Maheu, F. S. (2019). Chronic harsh parenting and anxiety associations with fear circuitry function in healthy adolescents: A preliminary study. *Biological Psychology*, 145, 198–210.
- Lange, N. (2012). Total and regional brain volumes in a population-based normative sample from 4 to 18 years: The NIH MRI study of normal brain development. *Cerebral Cortex*, 22, 1–12.
- Lenroot, R. K., & Giedd, J. N. (2006). Brain development in children and adolescents: Insights from anatomical magnetic resonance imaging. *Neuroscience and Biobehavioral Reviews*, 30, 718–729.
- LeWinn, K. Z., Sheridan, M. A., Keyes, K. M., Hamilton, A., & McLaughlin, K. A. (2017). Sample composition alters associations between age and brain structure. *Nature Communications*, 8, 874.
- Li, G., Wang, L., Yap, P.-T., Wang, F., Wu, Z., Meng, Y., ... Shen, D. (2019). Computational neuroanatomy of baby brains: A review. *NeuroImage*, 185, 906–925.
- Liberzon, I., Ma, S. T., Okada, G., Ho, S. S., Swain, J. E., & Evans, G. W. (2015). Childhood poverty and recruitment of adult emotion regulatory neurocircuitry. *Social Cognitive and Affective Neuroscience*, 10, 1596–1606. doi:10.1093/scan/nsv045
- Lim, L., Hart, H., Mehta, M., Worker, A., Simmons, A., Mirza, K., & Rubia, K. (2018). Grey matter volume and thickness abnormalities in young people with a history of childhood abuse. *Psychological Medicine*, 48(6), 1034–1046. doi:10.1017/S0033291717002392.
- Lim, L., Radua, J., & Rubia, K. (2014). Gray matter abnormalities in childhood maltreatment: A voxelwise meta analysis. *American Journal of Psychiatry*, 171, 854–863.
- Lupien, S. J., Leon, M. D., Santi, S. D., Convit, A., Tarshish, C., Thakur, M., ... Meaney, M. J. (1998). Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nature Neuroscience*, 1, 69–73.
- Marsh, R., Gerber, A. J., & Peterson, B. S. (2008). Neuroimaging studies of normal brain development and their relevance for understanding childhood neuropsychiatric disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*, 47, 1233–1251.
- Masten, A. S., & Cicchetti, D. (2010). Developmental cascades. *Development and Psychopathology*, 22, 491–495. doi:10.1017/S0954579410000222
- Mattison, R. E., & Spitznagel, E. L. (1999). Long-term stability of child behavior checklist profile types in a child psychiatric clinic population. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38, 700–707. doi:10.1097/00004583-199906000-00017
- McElroy, E., Belsky, J., Carragher, N., Fearon, P., & Patalay, P. (2018). Developmental stability of general and specific factors of psychopathology from early childhood to adolescence: Dynamic mutualism or p-differentiation? *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 59, 667–675. doi:10.1111/jcpp.12849
- McEwen, B. S., Nasca, C., & Gray, J. D. (2016). Stress effects on neuronal structure: Hippocampus, amygdala, and prefrontal cortex. *Neuropsychopharmacology*, 41, 3–23.
- McLaughlin, K. A., Peverill, M., Gold, A. L., Alves, S., & Sheridan, M. A. (2015). Child maltreatment and neural systems underlying emotion regulation. *Journal of the American Academy of Child and Adolescent Psychiatry*, 54(9), 753–762. doi:10.1016/j.jaac.2015.06.010.

- McLaughlin, K. A., Sheridan, M. A., Winter, W., Fox, N. A., Zeanah, C. H., & Nelson, C. A. (2014). Widespread reductions in cortical thickness following severe early-life deprivation: A neurodevelopmental pathway to attention-deficit/hyperactivity disorder. *Biological Psychiatry*, *76*, 629–638.
- McLaughlin, K. A., Weissman, D., & Bitrán, D. (2019). Childhood adversity and neural development: A systematic review. *Annual Review of Developmental Psychology*, *1*, 277–312.
- Mehta, M. A., Golemboski, N. I., Nosarti, C., Colvert, E., Mota, A., Williams, S. C. R., ... Sonuga-Barke, E. J. S. (2009). Amygdala, hippocampal and corpus callosum size following severe early institutional deprivation: The English and Romanian adoptees study pilot. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *50*, 943–951.
- Miller, G. A., & Chapman, J. P. (2001). Misunderstanding analysis of covariance. *Journal of Abnormal Psychology*, *110*, 40–48.
- Miller, D. J., Duka, T., Stimpson, C. D., Schapiro, S. J., Baze, W. B., McArthur, M. J., ... Sherwood, C. C. (2012). Prolonged myelination in human neocortical evolution. *Proceedings of the National Academy of Sciences*, *109*, 16480–16485.
- Mills, K. L., Goddings, A. L., Herting, M. M., Meuwese, R., Blakemore, S. J., Crone, E. A., ... Tammes, C. K. (2016). Structural brain development between childhood and adulthood: Convergence across four longitudinal samples. *NeuroImage*, *141*, 273–281.
- Molet, J., Maras, P. M., Kinney-Lang, E., Harris, N. G., Rashid, F., Ivy, A. S., ... Baram, T. Z. (2016). MRI uncovers disrupted hippocampal microstructure that underlies memory impairments after early-life adversity. *Hippocampus*, *26*, 1618–1632.
- Nanni, V., Uher, R., & Danese, A. (2012). Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: A meta-analysis. *American Journal of Psychiatry*, *169*, 141–151. doi:10.1176/appi.ajp.2011.11020335
- Neil, A. L., & Christensen, H. (2009). Efficacy and effectiveness of school-based prevention and early intervention programs for anxiety. *Clinical Psychology Review*, *29*, 208–215. doi:10.1016/j.cpr.2009.01.002
- Newman, L., Sivaratnam, C., & Komiti, A. (2015). Attachment and early brain development – neuroprotective interventions in infant–caregiver therapy. *Translational Developmental Psychiatry*, *3*, 28647.
- Newman, E., Thompson, W. K., Bartsch, H., Hagler, D. J., Chen, C. H., Brown, T. T., ... Jernigan, T. L. (2016). Anxiety is related to indices of cortical maturation in typically developing children and adolescents. *Brain Structure and Function*, *221*, 3013–3025. doi:10.1007/s00429-015-1085-9
- Panizzon, M. S., Fennema-Notestine, C., Eyler, L. T., Jernigan, T. L., Prom-Wormley, E., Neale, M., ... Kremen, W. S. (2009). Distinct genetic influences on cortical surface area and cortical thickness. *Cerebral Cortex*, *19*, 2728–2735. doi:10.1093/cercor/bhp026
- Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, *9*, 947–957.
- Prasad, K. M., Goradia, D., Eack, S., Rajagopalan, M., Nutche, J., Magge, T., ... Keshavan, M. S. (2010). Cortical surface characteristics among offspring of schizophrenia subjects. *Schizophrenia Research*, *116*, 143–151. doi:10.1016/j.schres.2009.11.003
- Qi, H., Ning, Y., Li, J., Guo, S., Chi, M., Gao, M., ... Wu, K. (2014). Gray matter volume abnormalities in depressive patients with and without anxiety disorders. *Medicine*, *93*, e345. doi:10.1097/MD.0000000000000345
- R Core Team. (2019). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. URL <https://www.R-project.org/>
- Rifkin-Graboi, A., Kong, L., Sim, L. W., Sanmugam, S., Broekman, B. F., Chen, H., ... Qiu, A. (2015). Maternal sensitivity, infant limbic structure volume and functional connectivity: A preliminary study. *Translational Psychiatry*, *5*, e668. doi:10.1038/tp.2015.133
- Rosso, I. M., Makris, N., Britton, J. C., Price, L. M., Gold, A. L., Zai, D., ... Rauch, S. L. (2010). Anxiety sensitivity correlates with two indices of right anterior insula structure in specific animal phobia. *Depression and Anxiety*, *27*, 1104–1110. doi:10.1002/da.20765
- Sapolsky, R. M. (1996). Stress, glucocorticoids, and damage to the nervous system: The current state of confusion. *Stress*, *1*, 1–19.
- Schwartz, C. E., Kunwar, P. S., Greve, D. N., Kagan, J., Snidman, N. C., & Bloch, R. B. (2012). A phenotype of early infancy predicts reactivity of the amygdala in male adults. *Molecular Psychiatry*, *17*, 1042–1050.
- Seligman, L. D., & Ollendick, T. H. (2011). Cognitive-behavioral therapy for anxiety disorders in youth. *Child and Adolescent Psychiatric Clinics of North America*, *20*, 217–238. doi:10.1016/j.chc.2011.01.003
- Semple, R. J., & Lee, J. (2007). *Mindfulness-based cognitive therapy for anxious children: A manual for treating childhood anxiety*. Oakland, CA: New Harbinger Publications.
- Smith, S. M., Nichols, T. E., Vidaurre, D., Winkler, A. M., Behrens, T. E., Glasser, M. F., ... Miller, K. L. (2015). A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nature Neuroscience*, *18*, 1565–1567. doi:10.1038/nn.4125
- Stiles, J., & Jernigan, T. L. (2010). The basics of brain development. *Neuropsychology Review*, *20*, 327–348.
- Strawn, J. R., Hamm, L., Fitzgerald, D. A., Fitzgerald, K. D., Monk, C. S., & Phan, K. L. (2015). Neurostructural abnormalities in pediatric anxiety disorders. *Journal of Anxiety Disorders*, *32*, 81–88.
- Strawn, J. R., John, W. C., Dominick, K. C., Swartz, M. S., Wehry, A. M., Patino, L. R., ... DelBello, M. P. (2014). Cortical surface anatomy in pediatric patients with generalized anxiety disorder. *Journal of Anxiety Disorders*, *28*, 717–723.
- Strawn, J. R., Wehry, A. M., Chu, W. J., Adler, C. M., Eliassen, J. C., Cerullo, M. A., ... DelBello, M. P. (2013). Neuroanatomic abnormalities in adolescents with generalized anxiety disorder: A voxel-based morphometry study. *Depression and Anxiety*, *30*, 842–848.
- Taylor, B. K., Eastman, J. A., Frenzel, M. R., Embury, C. M., Wang, Y. P., Stephen, J. M., ... Wilson, T. W. (2020). Subclinical anxiety and posttraumatic stress influence cortical thinning during adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*, S0890-8567(20)32216-4. Advance online publication. doi:10.1016/j.jaac.2020.11.020
- Tierney, A. L., & Nelson, C. A. (2009). Brain development and the role of experience in the early years. *Zero to Three*, *30*, 9–13.
- Tozzi, L., Garczarek, L., Janowitz, D., Stein, D. J., Wittfeld, K., Dobrowolny, H., ... for the ENIGMA-MDD Consortium. (2020). Interactive impact of childhood maltreatment, depression, and age on cortical brain structure: Mega-analytic findings from a large multi-site cohort. *Psychological Medicine*, *50*(6), 1020–1031. doi:10.1017/S003329171900093X.
- Valadez, E. A., Tottenham, N., Tabachnick, A. R., & Dozier, M. (2020). Early parenting intervention effects on brain responses to maternal cues among high-risk children. *The American Journal of Psychiatry*, *177*, 818–826. doi:10.1176/appi.ajp.2020.20010011
- VanTieghem, M., Korom, M., Flannery, J., Choy, T., Caldera, C., Humphreys, K. L., ... Tottenham, N. (2021). Longitudinal changes in amygdala, hippocampus and cortisol development following early caregiving adversity. *Developmental Cognitive Neuroscience*, *48*, 100916. doi:10.1016/j.dcn.2021.100916
- Verhulst, F. C., & Van der Ende, J. (1995). The eight-year stability of problem behavior in an epidemiologic sample. *Pediatric Research*, *38*, 612–617. doi:10.1203/00006450-199510000-00023
- Vidal-Pineiro, D., Parker, N., Shin, J., French, L., Jackowski, A. P., ... Mowinckel, A. M. (2020). Cellular correlates of cortical thinning throughout the lifespan. *Scientific Reports*, *10*, 21803.
- Vuoksimaa, E., Panizzon, M. S., Chen, C. H., Fiecas, M., Eyler, L. T., Fennema-Notestine, C., ... Kremen, W. S. (2015). The genetic association between neocortical volume and general cognitive ability is driven by global surface area rather than thickness. *Cerebral Cortex*, *25*, 2127–2137. doi:10.1093/cercor/bhu018
- Wiik, K. L., Loman, M. M., Van Ryzin, M. J., Armstrong, J. M., Essex, M. J., Pollak, S. D., & Gunnar, M. R. (2011). Behavioral and emotional symptoms of post-institutionalized children in middle childhood. *Journal of Child Psychology and Psychiatry*, *52*, 56–63.
- Winkler, A. M., Kochunov, P., Blangero, J., Almasy, L., Zilles, K., Fox, P. T., ... Glahn, D. C. (2010). Cortical thickness or grey matter volume? The importance of selecting the phenotype for imaging genetics studies. *NeuroImage*, *53*, 1135–1146. doi:10.1016/j.neuroimage.2009.12.028
- Winkler, A. M., Sabuncu, M. R., Yeo, B. T., Fischl, B., Greve, D. N., Kochunov, P., ... Glahn, D. C. (2012). Measuring and comparing brain cortical surface area and other areal quantities. *NeuroImage*, *61*, 1428–1443. doi:10.1016/j.neuroimage.2012.03.026
- Zhang, X., Luo, Q., Wang, S., Qiu, L., Pan, N., Kuang, W., ... Gong, Q. (2020). Dissociations in cortical thickness and surface area in non-comorbid never-treated patients with social anxiety disorder. *EBioMedicine*, *58*, 102910. doi:10.1016/j.ebiom.2020.102910